





## DRESSING OPTIONS

AN EXCLUSIVE COMBINATION OF  
TECHNOLOGY, SUPPORT AND PROVEN RESULTS



### V.A.C.® GranuFoam™ Dressing Promotes granulation

- Hydrophobic, open-pore structure facilitates exudate removal
- Provides an equal distribution of the negative pressure at the wound site
- Adapts to the contours of deep and irregular shaped wounds



### V.A.C.® WhiteFoam Dressing Provides comfort and versatility

- Recommended when granulation tissue needs to be controlled
- Higher tensile strength (than V.A.C.® GranuFoam™ Dressing) allows for easy removal from tunnels and undermining
- Helps promote flap and graft take
- Non-adherent material for a more comfortable dressing change



### V.A.C. GranuFoam Silver™ Dressing Helps protect from bacterial growth

- Rapid and effective microbial kill<sup>1</sup> based on *in-vitro* microbial testing
- Eliminates the need for adjunctive silver layers that may inhibit negative pressure and granulation
- Continuous exposure of silver ions directly to the wound bed
- Provides an effective barrier to bacterial penetration and may help reduce bioburden in wounds



<sup>1</sup>Based on *in-vitro* microbial ZOI Testing, KCI data on file.

**Note: Disposable components of the V.A.C.® Therapy System, including the foam dressings (V.A.C.® GranuFoam™, V.A.C. GranuFoam Silver™, or V.A.C.® WhiteFoam Dressing), are packaged sterile and are latex-free. All Dressings are for single use only. To help ensure safe and effective use, these named dressings are to be used only with V.A.C.® Therapy Units.**



An Acelity Company

**NOTE: Specific indications, contraindications, warnings, precautions and safety information exist for KCI products and therapies. Please consult a physician and product instructions for use prior to application. This material is intended for healthcare professionals only.**

©2011 KCI Licensing, Inc., All rights reserved. All trademarks designated herein, unless otherwise noted, are proprietary to KCI Licensing, Inc., and/or their respective affiliates and/or licensors. DSL#13-0601.CAE (Rev. 9/14)



Acelity™

## Dressing Selection

### Selecting an appropriate foam dressing

Wound Characteristics	V.A.C.® GranuFoam™ Dressing	V.A.C.® WhiteFoam Dressing	V.A.C. GranuFoam Silver™ Dressing	V.A.C.® GranuFoam™ Bridge Dressing
Deep, acute wounds with moderate granulation tissue present	•		•	
Full thickness pressure ulcers (Stage 3 or 4)	•		•	•
Flaps	•			
Painful wounds		•	•	
Superficial wounds		•		
Tunneling/sinus tracts/undermining		•		
Wounds that require controlled growth of granulation tissue		•		
Deep trauma wounds	•	•	•	
Diabetic foot ulcers	•	•	•	•
Dry wounds	•	•	•	
Post-graft placement (including dermal substitutes)	•	•		
Lower extremity ulcers, including Venous Leg Ulcers and Diabetic Foot Ulcers	•	•	•	•
Venous Insufficiency Ulcers	•	•	•	•
Need for barrier to bacterial penetration			•	

NOTE: These are general recommendations. As with any application, please consult the patient's treating physician about individual conditions and treatment.

Always consult the V.A.C.® Therapy Clinical Guidelines long with the applicable instructions for use, labeling and safety information before placing a V.A.C.® Therapy System on a patient.